

## *Supplementary Material*

# **Cognitive Change During the Life Course and Leukocyte Telomere Length in Late Middle-aged Men.**

PhD Lene Rask<sup>1,2,8,\*</sup>, PhD Laila Bendix<sup>3</sup>, PhD Maria Harbo<sup>4</sup>, PhD Birgitte Fagerlund<sup>5</sup>, Professor Erik Lykke Mortensen<sup>6,8</sup>, Professor Martin Lauritzen<sup>1,2,8,\*</sup>, Professor Merete Osler<sup>6,7</sup>

1. Department of Neuroscience and Pharmacology, University of Copenhagen, Copenhagen, Denmark
2. Department of Clinical Neurophysiology, Rigshospitalet - Glostrup, Glostrup, Denmark
3. Pain Research Group, Department of Anaesthesiology and Intensive Care Medicine, Odense University Hospital, Odense, Denmark
4. Department of Clinical Genetics, Vejle Hospital, Vejle, Denmark
5. Center for Neuropsychiatric Schizophrenia Research and Lundbeck Foundation Centre for Clinical Intervention and Neuropsychiatric Schizophrenia Research, University of Copenhagen, Psychiatric Centre Glostrup, Glostrup, Denmark.
6. Department of Public Health, University of Copenhagen, Copenhagen, Denmark
7. Research Center for Prevention and Health, Glostrup Hospital, Glostrup, Denmark
8. Center for Healthy Aging, University of Copenhagen, Copenhagen, Denmark

Corresponding authors: Lene Rask and Martin Lauritzen, [lene.rask.01@regionh.dk](mailto:lene.rask.01@regionh.dk) and [martin.johannes.lauritzen@regionh.dk](mailto:martin.johannes.lauritzen@regionh.dk)

**Supplementary Table 1** Studies of leukocyte telomere length (LTL) and cognitive decline

Martin-Ruiz 2006 [6]	MMSE	Non-demented	Stroke survivors	Cohort (2 y follow-up) LTL → Dementia	TRF	195	80±4y	Longer LTL associated with reduced risk of dementia and less reduction in MMSE	Age,BP,ApoE4, CVD
Yaffe 2009 [7]	MMSE, DSST	Non-demented	Health ABC	Cohort (7 y follow up) Cognitive decline → LTL	qPCR	2741	70-79y	Cognitive decline was less for longest tertile of LTL than medium and short for MMSE, not DSST	Age, gender, SEP
Mather 2010 [8]	Cognitive performance		Australian population	Cohort (4 y follow up) Cognitive decline → LTL	qPCR	646	40's & 60's	No cross-sectional association with LTL and cognitive function, nor to cognitive decline	Age, gender, smoking alcohol, exercise, BP
Devore 2011 [9]	TICS East Boston		Nurses Health study	Cohort (10y follow up) LTL → Cognitive decline	qPCR	2092	~65y ♀	Longer LTL associated with slower cognitive decline. Not a powerful marker.	Age, education, smoking, DM, BP
Harris 2012 [10]	MH-T, speed, WAIS-III	Relatively healthy	Lothian Birth Cohort	Cohort (60y follow up) Cognitive decline → LTL	qPCR	1048	70y	No association to cognitive performance at age 70 nor to change from 11-70y	Gender, education, smoking, alcohol

MMSE=Mini Mental State Examination; MHT=Moray house test; CANTAB=Cambridge Neuropsychological Test Automated Battery; DSST=Digit symbol substitution test; TICS=Telephone interview for cognitive status; WAIS=Wechsler Adult Intelligence Scale; TRF=Telomere restriction fragment length assay; BP= Blood pressure; BMI= Body Mass Index; SEP socioeconomic Position; CVD= Cardiovascular disease; DM= Diabetes; Baseline exposure → Outcome at follow-up

## References

1. Harris SE, Deary IJ, MacIntyre A, Lamb KJ, Radhakrishnan K, Starr JM et al. 2006 The association between telomere length, physical health, cognitive ageing and mortality in non-demented people. *Neuroscience Letters*; 406:260-4
2. Valdes AM, Deary IJ, Gardner J, Kimura M, Lu X, Spector TD et al. 2010 Leucocyte telomere length is associated with cognitive performance in healthy women. *Neurobiol Ageing*; 31:986-92
3. Bendix L, Gade MM, Staun PW, Kimura M, Jeune B, Hjelmborg JV et al. 2011 Leukocyte Telomere Length and Physical Ability among Danish Twins age 70+. *Mech Ageing Dev.* 132(11-12):568-72
4. Der G, Batty GD, Benzeval M, Deary IJ, Green MJ, McGlynn L et al. 2012 Is telomere length a biomarker for aging: cross-sectional evidence from the west of Scotland? *PLoS One.*;7:e45166
5. Ma SL, Lau ESS, Suen EW, Lam LC, Leung PC, Woo J et al. 2013 Telomere length and cognitive function in southern Chinese community-dwelling male elders. *Age & Ageing*; 42:450-5
6. Martin-Ruiz C, Dickinson HO, Keys B, Rowan E, Kenny RA, Von Zglinicki T et al. 2006 Telomere length predicts poststroke mortality, dementia, and cognitive decline. *Ann Neurol.*; 6:174-80
7. Yaffe K, Lindquist K, Kluse M, Cawthon R, Harris T, Hsueh WC et al. 2011 Telomere length and cognitive function in community-dwelling elders: Findings from the Health ABC study. *Neurobiology of ageing*; 32:2055-60

8. Mather KA, Jorm AF, Anstey KJ Milburn PJ, Easteal S, Christensen H. 2010 Cognitive performance and leukocyte telomere length in two narrow age cohorts: a population study. *BMC Geriatr.*; 10:62
9. Devore EE, Prescott J, De Vivo I, Grodstein F. 2011 Relative telomere length and cognitive decline in the Nurses' Health Study. *Neuroscience letter*; 492:15-8
10. Harris SE, Martin-Ruiz C, von Zglinicki T, Starr JM, Deary IJ. 2012 Telomere length and aging biomarkers in 70-year olds: the Lothian Birth Cohort 1936. *Neurobiol Aging*; 33:1486.e3-8